

42p

N 63 11008  
NASA TN D-1984

NASA TN D-1984

*Code 1*



# TECHNICAL NOTE

D-1984

PHOTOCHEMICAL PROBLEMS OF THE VENUS ATMOSPHERE

By

Paul Harteck, Robert R. Reeves, Jr.,  
and Barbara A. Thompson

Prepared under Grant No. NsG-261-62 by  
RENSSELAER POLYTECHNIC INSTITUTE  
Troy, New York  
for

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION  
WASHINGTON

June 1963

Code: 1

1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221  
2222  
2223  
2224  
2225  
2226  
2227  
2228  
2229  
2230  
2231  
2232  
2233  
2234  
2235  
2236  
2237  
2238  
2239  
2240  
2241  
2242  
2243  
2244  
2245  
2246  
2247  
2248  
2249  
2250  
2251  
2252  
2253  
2254  
2255  
2256  
2257  
2258  
2259  
2260  
2261  
2262  
2263  
2264  
2265  
2266  
2267  
2268  
2269  
2270  
2271  
2272  
2273  
2274  
2275  
2276  
2277  
2278  
2279  
2280  
2281  
2282  
2283  
2284  
2285  
2286  
2287  
2288  
2289  
2290  
2291  
2292  
2293  
2294  
2295  
2296  
2297  
2298  
2299  
2300  
2301  
2302  
2303  
2304  
2305  
2306  
2307  
2308  
2309  
2310  
2311  
2312  
2313  
2314  
2315  
2316  
2317  
2318  
2319  
2320  
2321  
2322  
2323  
2324  
2325  
2326  
2327  
2328  
2329  
2330  
2331  
2332  
2333  
2334  
2335  
2336  
2337  
2338  
2339  
2340  
2341  
2342  
2343  
2344  
2345  
2346  
2347  
2348  
2349  
2350  
2351  
2352  
2353  
2354  
2355  
2356  
2357  
2358  
2359  
2360  
2361  
2362  
2363  
2364  
2365  
2366  
2367  
2368  
2369  
2370  
2371  
2372  
2373  
2374  
2375  
2376  
2377  
2378  
2379  
2380  
2381  
2382  
2383  
2384  
2385  
2386  
2387  
2388  
2389  
2390  
2391  
2392  
2393  
2394  
2395  
2396  
2397  
2398  
2399  
2400  
2401  
2402  
2403  
2404  
2405  
2406  
2407  
2408  
2409  
2410  
2411  
2412  
2413  
2414  
2415  
2416  
2417  
2418  
2419  
2420  
2421  
2422  
2423  
2424  
2425  
2426  
2427  
2428  
2429  
2430  
2431  
2432  
2433  
2434  
2435  
2436  
2437  
2438  
2439  
2440  
2441  
2442  
2443  
2444  
2445  
2446  
2447  
2448  
2449  
2450  
2451  
2452  
2453  
2454  
2455  
2456  
2457  
2458  
2459  
2460  
2461  
2462  
2463  
2464  
2465  
2466  
2467  
2468  
2469  
2470  
2471  
2472  
2473  
2474  
2475  
2476  
2477  
2478  
2479  
2480  
2481  
2482  
2483  
2484  
2485  
2486  
2487  
2488  
2489  
2490  
2491  
2492  
2493  
2494  
2495  
2496  
2497  
2498  
2499  
2500  
2501  
2502  
2503  
2504  
2505  
2506  
2507  
2508  
2509  
2510  
2511  
2512  
2513  
2514  
2515  
2516  
2517  
2518  
2519  
2520  
2521  
2522  
2523  
2524  
2525  
2526  
2527  
2528  
2529  
2530  
2531  
2532  
2533  
2534  
2535  
2536  
2537  
2538  
2539  
2540  
2541  
2542  
2543  
2544  
2545  
2546  
2547  
2548  
2549  
2550  
2551  
2552  
2553  
2554  
2555  
2556  
2557  
2558  
2559  
2560  
2561  
2562  
2563  
2564  
2565  
2566  
2567  
2568  
2569  
2570  
2571  
2572  
2573  
2574  
2575  
2576  
2577  
2578  
2579  
2580  
2581  
2582  
2583  
2584  
2585  
2586  
2587  
2588  
2589  
2590  
2591  
2592  
2593  
2594  
2595  
2596  
2597  
2598  
2599  
2600  
2601  
2602  
2603  
2604  
2605  
2606  
2607  
2608  
2609  
2610  
2611  
2612  
2613  
2614  
2615  
2616  
2617  
2618  
2619  
2620  
2621  
2622  
2623  
2624  
2625  
26

## TABLE OF CONTENTS

	Page
SUMMARY	1
INTRODUCTION	3
EXPERIMENTAL	5
1. Absorption Spectroscopy - Absorption Coefficients of CO <sub>2</sub> , CO and O <sub>2</sub> between 1850 and 4000 Å	5
2. Irradiation Studies	11
KINETIC CONSIDERATIONS	19
MICROWAVE EMISSION	26
WATER AS A MINOR CONSTITUENT	31
THE CLOUD LAYER	32
CONCLUDING REMARKS	34
ACKNOWLEDGEMENTS	34
REFERENCES	36
TABLE	
I Abundance of Constituents of the Venus Atmosphere	4
II Radiation Chemistry of Carbon Dioxide	12
III Oxidation of CO by the O <sub>2</sub> - O <sub>3</sub> Cycle in the Lower Atmosphere	20
FIGURE	
1 Absorption Coefficient of Carbon Dioxide as a Function of Wavelength	7
2 Absorption Coefficient of Carbon Monoxide as a Function of Wavelength	8
3 Absorption Coefficient of Oxygen as a Function of Wavelength	9
4 Glow Discharge Through Carbon Dioxide in 22-Liter Flask	14
5 Dissociation of Carbon Dioxide by Ionizing Radiation as a Function of Pressure	16

FIGURE	Page
6 Absorption of Solar Radiation in the Venus Atmosphere	18
7 Atmospheric Characteristics for Earth and Venus	21
8 Experimental Arrangement for Studying Microwave Emission	30

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

---

TECHNICAL NOTE D-1984

---

PHOTOCHEMICAL PROBLEMS OF THE VENUS ATMOSPHERE

By

Paul Harteck, Robert R. Reeves, Jr.,  
and Barbara A. Thompson

SUMMARY

An experimental program to develop an understanding of the chemical processes occurring in the Venus atmosphere is in progress. Although the exact chemical composition of the Venus atmosphere is not known, it is known that  $\text{CO}_2$  is a major constituent,  $\text{CO}$  and  $\text{O}_2$  are minor components and there are probably substantial amounts of  $\text{N}_2$  and some noble gases. Water vapor seems to be almost entirely absent. At present there is no evidence for other gaseous constituents. In the Venus chemosphere  $\text{CO}_2$  will be dissociated into  $\text{CO}$  and  $\text{O}$ -atoms and at higher altitudes  $\text{N}_2$  and  $\text{CO}$  will be dissociated into atoms. These may recombine in several different ways. Experiments using both ultraviolet light and ionizing radiation have been carried out to supplement the information available in the literature on the mechanisms of these recombination reactions. Absorption coefficients of  $\text{CO}_2$ ,  $\text{CO}$ , and  $\text{O}_2$  have been determined in the region from 1850 to 4000 Å.

In connection with the recently publicized "high temperature" of Venus, it should be noted that there are a variety of alternate, nonthermal mechanisms to which the observed microwave

radiation can be attributed. The possibility of chemical origin of the microwave emission is discussed and preliminary experiments indicate that such an explanation may be feasible.

It is of interest that the atmosphere of Venus is chemically inert, being neither oxidizing nor reducing. This situation can be understood if Venus initially had a reducing atmosphere, as proposed by Urey and others, with excess CO. Under these conditions, over geologic ages, the CO would have been converted by the solar radiation to carbon suboxide polymer and to CO<sub>2</sub>. Thus, carbon suboxide polymer may cover the surface of Venus. Thermal decomposition of the suboxide may have occurred to a greater or lesser extent depending on the surface temperature. Such a polymer may be a major constituent of the visible clouds obscuring the surface from view. Even today carbon suboxide polymer may be forming and decomposing in an equilibrium steady state. Some other possible components of the cloud layer are suggested.

Since Venus apparently does not rotate, the dark side may be substantially cooler than the bright side; however, the difference may be less than anticipated from analogy with the Earth, because on the surface of a non-rotating planet, heat transfer by convection will be much more efficient than on a rotating planet. If the temperature on the dark side is very low (~234°K) and the partial pressure of CO<sub>2</sub> is about 10 atmospheres, as some have proposed, CO<sub>2</sub> may condense into liquid oceans.

## INTRODUCTION

The planet Venus is the nearest of the planets to Earth and in many ways can be regarded as a "twin" of Earth. Despite these facts, less is known about Venus than almost any other planet in our solar system, with the exception of Pluto. The principle reason for this mystery is the cloud layer covering the planet, which completely obscures the surface from view. Our direct knowledge of the nature of the temperature, pressure, chemical composition, and other conditions existing on Venus is confined to the region above the cloud layer. All information about the nature of the conditions at the surface of the planet must be derived from our knowledge of conditions above the cloud layer; the state of this knowledge is such that widely divergent views are possible.

It is the purpose of the work described in this paper to develop an understanding of the chemical processes occurring in the Venus atmosphere in the light of available data on atmospheric constituents, nature and abundance of impinging solar radiation, and the nature and rates of various possible competing chemical reactions.

The first information about the constitution of the Venus atmosphere came in 1932 when Adams and Dunham detected strong absorption bands of carbon dioxide (for discussion see Dunham, 1952). Since that time a number of workers have searched unsuccessfully for other constituents. In the past year the presence of traces of oxygen has been reported by Prokofiev and Petrova (1962), and Sinton (1962) has observed a small amount of carbon monoxide. Traces of water vapor have been reported, but this result is uncertain because it is based on one observation from a balloon without precise correction for water vapor in the Earth's atmosphere (Strong, 1961). These data are summarized in Table I. It is striking that oxygen and carbon monoxide, the decomposition products of carbon dioxide, are present in such small amounts. It should be noted that nitrogen is expected to be present in relatively large amounts because of its high cosmic abundance, since

Table I

Abundance of Constituents of the Venus Atmosphere

<u>Constituent</u>	<u>Abundance above Cloud Layer</u>	<u>Reference</u>
CO <sub>2</sub>	10 <sup>5</sup> cm atm	Dunham (1952), Herzberg (1952)
O <sub>2</sub>	(a) < 100 cm atm (b) detected	Dunham (1952) Prokofiev and Petrova (1962)
CO	(a) < 100 cm atm (b) a few cm atm	Kuiper (1952) Sinton (1962)
H <sub>2</sub> O	~ 2.5 cm atm	Strong (1961)
N <sub>2</sub> O	< 100 cm atm	Kuiper (1952)
NH <sub>3</sub>	< 4 cm atm	Kuiper (1952)
CH <sub>4</sub>	< 20 cm atm	Kuiper (1952)
C <sub>2</sub> H <sub>4</sub>	< 3 cm atm	Kuiper (1952)
C <sub>2</sub> H <sub>6</sub>	< 1 cm atm	Kuiper (1952)

it can neither escape from the gravitational field of Venus nor be consumed chemically. However, it is extremely difficult to detect by Earth-based measurements because of the strong absorption of the Earth's atmosphere in the far ultraviolet region where nitrogen absorbs. A future Mariner type probe may certainly carry suitable instrumentation for nitrogen detection and overcome this problem. Similar considerations apply to the rare gases, which should be expected to be present in abundances comparable to those in the Earth's atmosphere. However, the escape conditions for helium-3 and helium-4 would be more favorable on Venus than on Earth. Some efforts have been made to observe emission from  $N_2^+$  in the night sky of Venus, but these have yielded somewhat uncertain results( Kozyrev, 1961; Newkirk, 1961).

To provide an additional basis for detailed discussion of the chemistry of Venus atmosphere, certain useful data can be obtained by laboratory experiments. The results of these experiments are described in the following section.

#### EXPERIMENTAL

##### 1. Absorption Spectroscopy-Absorption Coefficients of $CO_2$ , CO and $O_2$ between 1850 and 4000 A.

In considering the photochemistry of the Venus atmosphere the two things which must be first considered are the intensities of the various solar radiations impinging on the atmosphere and the absorption coefficients of the various atmospheric constituents for these radiations. The solar radiation intensities are well known for Earth (Johnson, 1961) and can be readily calculated for Venus by the inverse square law. Since the distance of Venus from the sun is about 0.7 that for the distance of the Earth from the sun, Venus receives about twice as much solar radiation as the Earth. Carbon dioxide, the major constituent\*, is known to absorb fairly strongly below 1700 A

\*If it should be determined in the future that  $N_2$  is more abundant than  $CO_2$ , the kinetics of the chemistry of the Venus atmosphere would remain virtually unchanged.

with a maximum absorption coefficient of about  $30 \text{ cm}^{-1}$  at  $\sim 1450 \text{ \AA}$ , corresponding to dissociation into CO and O ( $^1\text{D}$ ). Absorption at wavelengths between 1700 and 2250  $\text{\AA}$ , corresponding to dissociation into CO and O in their ground states has not been reported. However, this region is of paramount importance for the photochemistry of the Venus atmosphere, since even if the dissociation occurs with only a very small absorption coefficient ( $10^{-4} \text{ cm}^{-1}$  or less), the increased solar radiation intensity at these wavelengths and the long path length through  $\text{CO}_2$  would make it a very significant process. A knowledge of the magnitudes of the absorption coefficients of the minor constituents CO and  $\text{O}_2$  in this wavelength region is similarly of great interest. To extend the information available in the literature, measurements also were made on these gases in the course of the present study. A Perkin-Elmer model 350 absorption spectrophotometer which covers the entire ultraviolet, visible, and near infrared regions with a short wavelength limit of about 1850  $\text{\AA}$  was used. Absorption coefficients as low as  $10^{-4} \text{ cm}^{-1}$  can be determined with this instrument. We have measured absorption spectra of all the gases of interest for the Venus atmosphere in the region between 1850 and 4000  $\text{\AA}$ . This supplements the data of Watanabe et al. (1953) whose results in general cover wavelengths below 1800  $\text{\AA}$ . The gases measured included  $\text{CO}_2$ , CO,  $\text{O}_2$ ,  $\text{CH}_4$ ,  $\text{H}_2\text{O}$ ,  $\text{N}_2\text{O}$ , NO,  $\text{NO}_2$ ,  $\text{SO}_2$ , and  $\text{NH}_3^*$ . Most of the gases were carefully prepurified to remove traces of impurities which might cause misleading results. This was particularly necessary in the case of carbon monoxide since even the research grade gas (obtained from Air Reduction Co.) contained several hundred ppm of iron carbonyl which absorbs strongly in the ultraviolet. The results obtained for  $\text{CO}_2$ , CO and  $\text{O}_2$  are shown in Figures 1, 2, and 3.

---

\*Not all of these measurements are reported in this paper.




Figure 1 ABSORPTION COEFFICIENT OF CARBON DIOXIDE AS A FUNCTION OF WAVELENGTH




Figure 2 ABSORPTION COEFFICIENT OF CARBON MONOXIDE AS A FUNCTION OF WAVELENGTH




Figure 3 ABSORPTION COEFFICIENT OF OXYGEN AS A FUNCTION OF WAVELENGTH

### CO<sub>2</sub> (Figure 1)

It was found that CO<sub>2</sub> begins to absorb at about 1960 Å and gradually increases absorption with decreasing wavelengths. Between 1960 Å and 4000 Å no absorption could be detected, thus indicating an absorption coefficient of less than  $10^{-4} \text{ cm}^{-1}$  in this region.

### CO (Figure 2)

Carbon monoxide is the gas with the highest binding energy known. Its dissociation energy of 11.11 eV corresponds to a wavelength of 1110 Å. Thus no continuous absorption is to be expected in the entire range above 1110 Å. The wavelengths which dissociate CO are energetic enough to ionize NO and are in the region normally referred to as ionizing radiation. In our work with CO we could observe the absorption peaks of the Cameron bands, representing the forbidden transition  $X^1\Sigma \rightarrow a^3\Pi$ , in the region between 1850 and 2060 Å. It should be noted that the absorption coefficients for these bands are averages since the closely packed rotational structure within the bands is not resolved. Although these average absorption coefficients are rather low, it should be noted that if as much as 10 cm-atm of CO is present in the Venus atmosphere, about 3% of the impinging light quanta in the wavelength interval from 1850 to 2060 Å could be absorbed. Since in this wavelength interval about  $10^{13}$  light quanta per  $\text{cm}^2\text{-sec}$  are impinging on the atmosphere, as many as  $3 \times 10^{11}$  absorption processes may occur resulting in CO in the metastable  $a^3\Pi$  state. The chemical reactions of this CO excited state are not known at present and constitute a rather interesting field of research.

### O<sub>2</sub> (Figure 3)

In the case of oxygen, the data of Watanabe, et al. (1953) have been extended here to include the 0,0; 1, 0; 2, 0; and 3,0 Schumann-Runge bands in the region from 1920 to 2020 Å. In addition it was determined that the absorption coefficient in the wavelength

region above 2100 Å must be less than  $10^{-4} \text{ cm}^{-1}$  since we were unable to detect any absorption.

## 2. Irradiation Studies

### A. Ionizing Radiation

It is well known that carbon dioxide at atmospheric pressure is very resistant to decomposition by ionizing radiation (Harteck and Dondes, 1955; 1957; Lind, 1961), the radiation equilibrium with ionizing radiation being less than 0.1% decomposed\*. The explanation proposed for this high stability involves regeneration via a carbon suboxide intermediate (see Table II). In the presence of an inhibitor, however, carbon dioxide becomes decomposed by ionizing radiation. An excellent inhibitor is  $\text{NO}_2$ ; less effective inhibitors include  $\text{SO}_2$  and  $\text{I}_2$ . It is not expected that  $\text{NO}_2$  is present in the Venus atmosphere in view of the results obtained by Harteck (1957) and Dondes (1957) who studied the stationary concentrations of  $\text{NO}$  and  $\text{NO}_2$  in a mixture of  $\text{O}_2$  and  $\text{N}_2$  subjected to ionizing radiation at various pressures. Extrapolation to low pressures shows that almost no  $\text{NO}$  or  $\text{NO}_2$  is present. Harteck and Dondes (1958) have also investigated mixtures of  $\text{CO}_2$  and  $\text{N}_2$  at high pressures and find that since N-atoms do not react directly with  $\text{CO}_2$ , the stationary concentration of  $\text{NO}$ ,  $\text{NO}_2$  and  $\text{N}_2\text{O}$  is always less in a mixture of  $\text{CO}_2$ ,  $\text{N}_2$ , and  $\text{O}_2$  than in a mixture containing only  $\text{N}_2$  and  $\text{O}_2$ . Therefore,  $\text{NO}$  and  $\text{NO}_2$  should be absent from the entire atmosphere of Venus since the concentration must be less than that in the Earth's atmosphere where oxides of nitrogen play no role except as  $\text{NO}^+$  in the ionospheric layers.

Carbon dioxide behaves in an entirely different manner with ultraviolet radiation such as the Xenon resonance line of 1470 Å. As Groth (1937) working at atmospheric pressure, and Mahan (1960) and

---

\*Actually, under very clean conditions there is practically no stationary state of  $\text{CO}$  and  $\text{O}_2$  as dissociation products, but since  $\text{CO}$  is polymerized to carbon suboxide in the gas phase a small amount of oxygen will remain from this process to inhibit further formation of suboxide.

Table II

Radiation Chemistry of Carbon Dioxide

(Harteck and Dondes, 1955)



Jucker and Rideal (1957) at lower pressures, have shown,  $\text{CO}_2$  becomes dissociated with a quantum yield of one. Since the solar radiation is a mixture of ionizing and ultraviolet radiation, it was necessary to make additional experiments. Initial experiments were carried out to investigate the effect of ionizing radiation at lower pressures on  $\text{CO}_2$ . It was convenient to do this with a glow discharge. As is well known, the effect of a glow discharge at low pressures is comparable to that of other modes of ionizing radiation.

The apparatus used consisted of a 22-liter Pyrex flask into which two aluminum electrodes were sealed, as shown in Figure 4. Both AC and DC discharges were employed, the power level in all cases being about 0.6 watts distributed over the vessel area. This power level was a compromise between the intensity of solar ionizing radiation and the need to reduce the time of experiments to a reasonable value. Under these experimental conditions the radiation intensity was about five times that of the ionizing radiation from the sun. The solar ionizing radiation intensity can be considered to be the total energy input for light quanta of wavelengths shorter than 1100-1200 Å. This total ionizing radiation is equivalent to about  $10^{12}$  ionization processes per  $\text{cm}^2\text{-sec}$  in the Venus atmosphere. Experiments were carried out at two initial  $\text{CO}_2$  pressures, 500 $\mu$  and 250 $\mu$ . At the conclusion of each experiment the reaction mixture was passed through two traps cooled with liquid nitrogen, the second of which contained activated charcoal. Undecomposed  $\text{CO}_2$  was frozen out in the first trap and dissociation products uncondensed at liquid nitrogen temperature were adsorbed on the charcoal. The final  $\text{CO}_2$  pressure was determined and used as a measure of the extent of decomposition. The products adsorbed on the charcoal were analyzed mass spectrometrically and in all cases were found to be CO and  $\text{O}_2$  in approximately the stoichiometric 2:1 ratio. At both pressures a rapid initial decomposition was observed which decreased as equilibrium was approached. At equilibrium the fraction decomposed was about 0.37 for 500 $\mu$  initial pressure




Figure 4 GLOW DISCHARGE THROUGH CARBON DIOXIDE IN 22-LITER  
FLASK

and about 0.50 for 250 $\mu$  initial pressure. The presence of excess CO or O<sub>2</sub> initially did not change the position of the equilibrium significantly.

The chemosphere of Venus will exist, however, in a pressure range similar to that of the Earth, i.e., in a pressure range of a few microns. In connection with experiments described below, extrapolation can be made to the pressure range of microns or less showing that a quantitative dissociation of CO<sub>2</sub> and O should prevail.

To obtain experimental data over a larger pressure range and to determine whether the mode of ionizing radiation is important for the decomposition of CO<sub>2</sub>, a series of samples was irradiated with mixed reactor irradiation ( $\beta$ ,  $\gamma$ , and ionizing radiation caused by collisions with fast neutrons) in the Brookhaven reactor. The conditions were nearly identical to those used by Harteck and Dondes (1955, 1957) who irradiated CO<sub>2</sub> at higher pressures. The CO<sub>2</sub> pressure of these irradiated samples was varied from 250 mm down to 2.5 mm. The irradiations were carried out at room temperature which may not be too different from that of the Venus chemosphere. The time of the irradiations was sufficiently long that equilibrium should have been reached. After irradiation the CO<sub>2</sub> was frozen out and the pressure of the decomposition products was measured. These were then adsorbed onto charcoal and analyzed mass spectrometrically. The CO<sub>2</sub> pressure was then measured and was also mass analyzed, to check for impurities. Figure 5 shows a logarithmic plot of the ratio of CO<sub>2</sub> dissociated to CO<sub>2</sub> undissociated, as a function of pressure, including both the glow discharge and reactor irradiations. It is clear that dissociation increases rapidly with decreasing pressure and that the type of ionizing radiation is not important.

In addition a few experiments were made with mixtures of nitrogen and CO<sub>2</sub>. As expected, there was no difference in the results because as discussed above, under these conditions the stationary concentration of NO<sub>2</sub>, which may act as an inhibitor, is so low that no




Figure 5 DISSOCIATION OF CARBON DIOXIDE BY IONIZING RADIATION AS A FUNCTION OF PRESSURE

increased dissociation occurs. Therefore, even if substantial amounts of  $N_2$  are present in the Venus atmosphere, these will not affect the photochemistry.

#### B. Mixed Ultraviolet Radiation

The ultraviolet spectrum can be divided into two regions for which the photochemical behavior of the Venus atmosphere will be entirely different due to the nature of the absorption coefficients of  $CO_2$  and  $O_2$ . In the region of wavelengths below about 1900 Å, both  $CO_2$  and  $O_2$  will absorb with high absorption coefficients so that all of this ultraviolet radiation is absorbed at a pressure of a few microns. (See Figure 6). In this region, which has been very thoroughly investigated from the standpoint of the behavior of oxygen in the Earth's atmosphere, oxygen is almost entirely dissociated ( $N_2$  does not absorb at all) and  $CO_2$  will also be largely dissociated although its absorption coefficient is only about 0.1 that of  $O_2$  (See Figures 1 and 3). Radiation in this region is thus responsible for the chemosphere which is a region of great interest.

At longer wavelengths the absorption coefficients of both  $CO_2$  and  $O_2$  are quite small and above 2250 Å  $CO_2$  cannot dissociate. Weak absorption by oxygen at these wavelengths can result in the formation of  $O_3$ , however, which absorbs strongly, regenerating oxygen atoms. This region is thus of great interest from the standpoint of determining whether the carbon monoxide formed in the chemosphere of Venus has a rather long lifetime in the lower levels of the Venus atmosphere or whether it is oxidized rapidly.

To answer this question, we have subjected pure  $CO_2$  with a few percent argon added, and mixtures of 90%  $CO_2$ , 7% Ar, 2%  $O_2$ , and 1% CO to irradiation from a 500 watt, high pressure mercury lamp. The argon addition was made for mass spectrometer standardization purposes. The radiation from this lamp was analyzed with a Jarrell-Ash recording monochromator and it was found that the 2537 Å line was almost completely self-reversed. The shortest line of high intensity was at 2253 Å





Figure 6 ABSORPTION OF SOLAR RADIATION IN THE VENUS ATMOSPHERE

wavelength since shorter wavelengths were absorbed by the cooling water jacket. The sample pressure in each case was about 500 mm. During each irradiation small samples were withdrawn at intervals, the CO<sub>2</sub> in each was frozen out with liquid nitrogen, and the uncondensed residue analyzed with the mass spectrometer. By recording the ratio of the peak heights at mass 40 to those at masses 32 and 28 any change in the CO and O<sub>2</sub> could be detected. It was found that both the CO and O<sub>2</sub> concentrations decreased with time and it was estimated that for each light quantum primarily absorbed by O<sub>2</sub>, about one or two CO molecules become oxidized. With pure CO<sub>2</sub>, no change was detected after irradiation for 24 hours confirming that no radiation was present sufficiently energetic to dissociate CO<sub>2</sub>.

Under the same conditions mixtures of Ar, CO and O<sub>2</sub> but without CO<sub>2</sub>, were irradiated and gave about the same rate of oxidation, about 1% per hour. Thus it appears that excited oxygen does not become deactivated by the presence of abundant CO<sub>2</sub>, and the reverse reaction, which could have been postulated:



obviously does not occur.

When pure oxygen was irradiated, it was observed that a substantial equilibrium concentration of ozone was built up (4 mg/l.). When CO was present, practically no stationary ozone could be observed. This indicates that CO is oxidized to CO<sub>2</sub> via the oxygen-ozone cycle as shown in Table III.

#### KINETIC CONSIDERATIONS

On the basis of the experimental data certain conclusions can now be drawn about the state of the Venus atmosphere as a function of altitude, making comparisons with the Earth's atmosphere, under the assumption that no other chemical species, at present unknown, are present in the gas phase. The comparison is shown schematically in Figure 7.

Table III

Oxidation of CO by the  $O_2 - O_3$

Cycle in the Lower Atmosphere






Figure 7 ATMOSPHERIC CHARACTERISTICS FOR EARTH AND VENUS

In the atmosphere there are three or four distinct levels. The highest is the exosphere from which molecules escape without collisions. Since Venus is very dry, no H-atoms formed by photodissociation of water vapor will be present in this region. However, a stationary H-atom concentration will exist due to the solar winds. This can be estimated as about 1/100 or 1/1000 of the H-atom concentration in the Earth's atmosphere. Since at 100 km (or  $10^{-6}$  atm) in the Earth's atmosphere there are  $3 \times 10^7$  H-atoms/cc, there may be about  $10^5$  H-atoms/cc at a corresponding altitude on Venus.

Below the exosphere is the ionosphere. Since the major constituent of the Venus atmosphere is  $\text{CO}_2$ , the Venus ionosphere will differ from that of Earth. High percentages of ions such as  $\text{CO}_2^+$  and  $\text{CO}^+$  may be expected. Other more complex ions may also be present.

The third region, the chemosphere, is the one in which the far ultraviolet radiation from about 1200 to 1900 Å is strongly absorbed; above 1900 Å the absorption coefficients for  $\text{CO}_2$ ,  $\text{O}_2$ , and CO become small. In the Earth's atmosphere in this region solar ultraviolet radiation dissociates oxygen into oxygen atoms. Above 100 kilometers altitude most of the oxygen is in the atomic form. Recombination occurs mainly by three-body collisions such as



followed by



At radiation equilibrium the total dissociation by light absorption must equal the total recombination:

$$\Sigma h\nu = \Sigma \text{ recombination} \quad (12)$$

The recombination rate is third order and therefore increases approximately as the third power of the pressure. On the basis of these facts

the altitude and pressure can be calculated at which the chemosphere begins. It may be emphasized that  $N_2$  plays only a minor part in the chemosphere since its absorption coefficient is significant only in the region of 800 Å wavelength and below.\*

Similar conditions for a chemosphere will prevail on Venus. If  $CO_2$  alone were present then dissociation would occur predominantly to CO and O-atoms and the following two competing recombination reactions would occur:



Reaction (6) is slow, however, by comparison to Reaction (11) by a factor of about 1000 at room temperature. This ratio will be somewhat temperature dependent because Reaction (6) has a heat of activation of several kcal. The situation in the Venus chemosphere is more complicated however, because molecular oxygen is formed by Reaction (11) and has a higher absorption coefficient than  $CO_2$ . Also, as seen in Figure 3, this absorption of  $O_2$  extends somewhat further into the longer wavelength region.

The relative concentrations of  $CO_2$ , CO,  $O_2$ , and O-atoms in the Venus chemosphere and above can be estimated in the following way. In this region the rates of decomposition of  $CO_2$  and  $O_2$  which may be present are:

$$\frac{-d(CO_2)}{dt} = I \alpha^I(CO_2) \quad (13)$$

$$\frac{-d(O_2)}{dt} = I \alpha^{II}(O_2) \quad (14)$$

where I is the number of impinging light quanta capable of producing dissociation and  $\alpha^I$  and  $\alpha^{II}$  are the absorption coefficients for  $CO_2$

---

\*Kinetic data for the reactions of interest in the chemosphere have been presented in detail by Harbeck and Reeves (1962).

and  $O_2$  respectively. Similarly, the rates of formation of  $CO_2$  and  $O_2$  by three-body recombination\* are given by:

$$+ \frac{d(CO_2)}{dt} = k^I (CO)(O)(M) \quad (15)$$

$$+ \frac{d(O_2)}{dt} = k^{II}(O)(O)(M) \quad (16)$$

where  $k^I$  and  $k^{II}$  are the respective three-body rate constants. At equilibrium the rate of formation will be equal to the rate of decomposition for each component and therefore the following relationship will hold:

$$\frac{-d(CO_2)/dt}{-d(O_2)/dt} = \frac{+d(CO_2)/dt}{+d(O_2)/dt} \quad (17)$$

or

$$\frac{\alpha^I(CO_2)}{\alpha^{II}(O_2)} = \frac{k^I(CO)}{k^{II}(O)} \quad (18)$$

The ratio  $\alpha^I/\alpha^{II}$  is about 1/20 over the wavelength region of interest and the ratio  $k^I/k^{II}$  is about 1/1000, as noted above. Thus:

$$\frac{(CO_2)}{(CO)} \times \frac{(O)}{(O_2)} \approx \frac{1}{50} \quad (19)$$

From this is immediately apparent that in the region where the total pressure is about  $10^{-7}$  atmospheres and the ratio  $(O)/(O_2)$  is about 10, the ratio  $(CO_2)/(CO)$  will be about 1/500. It should be noted that the entire atmosphere above the chemosphere is equivalent to only about 1 cm-atm so that these considerations do not imply the presence of extensive amounts of atmospheric CO and  $O_2$ . As the pressure increases other recombination mechanisms will become significant and the above relationship will no longer be valid.

\*For simplicity we have assumed that three-body recombination predominates for  $CO_2$  formation; however, it should be kept in mind that at low pressures two-body recombination may become important.

The molecular oxygen formed in the chemosphere will be carried to lower regions of the atmosphere by diffusion and convection. Since it is known that oxygen is only a minor constituent of the Venus atmosphere, it cannot be continually built up by this photochemical process. Reoxidation of the carbon monoxide must therefore occur via the ozone cycle at these levels below the chemosphere as mentioned previously. As seen from Table III the O-atoms formed by Reaction (7) may simply cycle over the oxygen-ozone cycle:



until reaction with CO occurs forming back CO<sub>2</sub> (Reaction (6)). The ozone molecule has a strong absorption in the ultraviolet region to almost 3000 Å and will be dissociated within a few seconds while recombination to ozone will depend on the total pressure and oxygen concentration and will vary from hours to fractions of a second. The extent to which reaction (9) occurs will depend upon the relative concentrations of O<sub>3</sub> and CO. The absolute concentrations of all constituents in the regions below the chemosphere will depend on diffusion and convection conditions which are at present unknown.

Carbon monoxide will become excited and dissociated by ultraviolet light in the upper atmosphere. It can be estimated that  $10^{11} - 10^{12}$  carbon atoms per cm<sup>2</sup>-sec are formed by ionizing radiation (assuming  $10^{12}$  ionizing quanta/cm<sup>2</sup>-sec with a M/N value of ~ 1 for the decomposition of CO). These carbon atoms will obviously recombine with O-atoms forming CO in three-body collisions.\* Similarly, reaction with CO to form carbon suboxide may occur:



Both reactions are third order and probably of similar rate.\*\* As has

\*Reactions of carbon atoms with various gases have recently been studied by Dubrin, MacKay, and Wolfgang (1963).

\*\*The reaction of carbon atoms with the small amounts of molecular oxygen present should be negligible, especially since the rate is slow (von Weyssenhoff, Dondès, and Harteck, 1962).

been shown by von Weyssenhoff, Dondes and Harteck (1962) the reaction of carbon suboxide with O-atoms is exceedingly fast giving both




and



Thus it appears that under existing conditions there is not much possibility that carbon suboxide accumulates in the Venus chemosphere. However, this statement deserves further critical consideration since many data are not well enough known at present.

In addition, it has been shown (Lind, 1961; von Weyssenhoff, Harteck, and Dondes, 1963) that excited CO molecules can react with CO to give carbon suboxide or CO<sub>2</sub>:



These reactions may also occur to a certain extent in the upper atmosphere of Venus. Since the concentration of CO is low, there may not be an opportunity for an excited CO molecule to react with a second CO molecule, but alternate reactions may occur, such as with O<sub>2</sub> to form CO<sub>2</sub> + O.

In the upper atmosphere of Venus nitrogen will be dissociated into N-atoms and these may react with the carbon atoms produced by dissociation of CO to form CN or cyanogen, C<sub>2</sub>N<sub>2</sub>. Under the action of ionizing radiation cyanogen may polymerize to (CN)<sub>x</sub> or, in the presence of oxygen, to (CNO)<sub>x</sub>, a pale yellow solid (Lind, 1961). The extent to which such reactions occur in the Venus atmosphere is not now known and the possibility that such polymers may contribute to the observed clouds should be considered.

#### MICROWAVE EMISSION

In the last few years radio astronomers have observed the emission of radiation in the microwave region ( $\lambda = 0.4 - 10.3$  cm) from

Venus (see, for example, Mayer, 1961 and Kuz'min and Salomonovich, 1962). This emission has been confirmed by the recent flight of Mariner II (Jet Propulsion Laboratory, 1963). The explanations which have been advanced to explain the origin of this radiation can be divided into two groups according to whether the origin is considered to be thermal (Sagan, 1960; Opik, 1961) or non-thermal (Jones, 1961; Tolbert and Straiton, 1962; Scarf, 1963). The most popular current explanation assumes that the microwave radiation is thermal in origin and corresponds to a high surface temperature for the planet. In particular, it is proposed that a high temperature ( $\sim 700^{\circ}\text{K}$ ) does exist at the surface with very cold clouds ( $\sim 230^{\circ}\text{K}$ ) above (Sagan, 1961; Jet Propulsion Laboratory, 1963; see also Kellogg and Sagan, 1961).

Prior to the discovery of the microwave emission, estimates of the temperature of Venus had been made based on a knowledge of the heat flux from the sun, the albedo of Venus, and estimates of the heat balance. The resulting temperature was found to be similar to that of Earth (Kuiper, 1952). A somewhat higher temperature might be estimated from our current knowledge that the planet rotates extremely slowly, if at all.

With the reports of emission of microwave radiation calculations of the temperature required for Venus to emit this radiation as a blackbody radiator have been made and it is from these calculations that the reports of  $600^{\circ}\text{--}700^{\circ}\text{K}$  are derived. No other evidence for such high temperatures is known and all current models which postulate an insulating atmospheric blanket to justify such high surface temperatures seem more sophisticated than realistic. The absence of water vapor and the presence of very cold clouds constitute major stumbling-blocks to the acceptance of any "greenhouse" theory involving surface temperatures of the order of  $700^{\circ}\text{K}$  (Sagan, 1961; Kellogg and Sagan, 1961). It may be noted that Chamberlain and Kuiper (1956) have calculated a temperature of  $285^{\circ}\text{K}$  from the rotational structure of the 7820 Å and

8689 Å bands in the CO<sub>2</sub> absorption spectrum of the Venus atmosphere. Spinrad (1962) has recently made a new calculation using the 7820 Å band structure as recorded on plates obtained by Adams and Dunham (1932) and making slightly different assumptions. Spinrad's calculated temperatures range from 214°K to 445°K with the extreme values observed for the same phase angle. As Chamberlain and Kuiper (1956) have pointed out, the accuracy of an individual measurement of this type is rather low. Thus, the overall spread most likely results from errors inherent in the measurement rather than from true variations in the CO<sub>2</sub> temperature existing in the Venus atmosphere. Therefore, the average of all the values, 337°K, should be much less subject to error than the extremes, e.g., the 445°K emphasized by Spinrad.

Because of these fundamental difficulties involved in a thermal origin of the microwave emission and because the surface temperature is obviously of great importance for chemical and biological considerations, various non-thermal origins of the observed microwave radiation have been proposed (Jones, 1961; Tolbert and Straiton, 1962; Scarf, 1963). However, these non-thermal models again have various drawbacks. Thus, it is appropriate to seek alternate sources for the microwave emission, and, as chemists, we have investigated the possibility of a chemical origin.

It is well known that the reaction of carbon monoxide with oxygen may result in emission of light both in form of bands and a strong continuum (See Gaydon, 1957). This reaction certainly occurs to a significant extent on Venus. The continuum is believed to arise from the reaction



Clearly, if only the spectrum of this radiation were considered in deriving a temperature for the reaction, an unreasonably high temperature would be obtained. Thus, an unexpectedly high emission in a certain spectral region should not be considered as black-body radiation

from the actual thermal temperature could be derived\*. It must be kept in mind that a system in which chemical reactions occur is a system which is not in thermodynamic equilibrium.

Gaydon (1957 ; 1961) has proposed that the continuum emitted in the carbon monoxide-oxygen reaction occurs as the CO and O approach each other on a repulsive potential curve and drop down to the stable ground state. The continuum has been observed to extend to wavelengths as long as 9000 Å (Feast, 1950) and there seems every reason to expect it to extend to much longer wavelengths. The question thus arises as to whether this emission extends into the microwave region, thus providing a possible chemical origin for the observed microwave emission from Venus. In addition, the possibility of other chemical reactions of a similar nature which may result in microwave emission should be considered. We now have in progress a survey program to evaluate the possibility of chemical reactions as a source of microwave emission.

Through the courtesy of the Naco Electronics Corporation we have been able to analyze the radiation from the various reactions for emission in the microwave region. As a matter of convenience we chose to examine the wavelength region between 2.8 and 4.1 cm (7.4 - 12 kMc/s known as X-band. The apparatus is shown in Figure 8. The system used had a lower detection limit of about  $10^{-11}$  watts with about a 10 micron window. To date it has not been possible to observe any signal from reactions between atoms and molecules. However, from the discharge itself, signals were detected throughout the entire X-band using CO<sub>2</sub> or SO<sub>2</sub>, although most gases such as air gave no signal. The intensity of this microwave emission corresponds to an unrealistically high blackbody temperature, in excess of 100,000°K. It therefore seems probable that the Venus microwave emission may be caused, at least in

---

\*The "high" microwave emission observed at 10 cm wavelength would represent only  $10^{-16}$  of the maximum intensity expected from a blackbody radiator at a temperature of 600°K ( $\lambda_{\text{max}} = 4.9 \mu$ )





Figure 8 EXPERIMENTAL ARRANGEMENT FOR STUDYING MICROWAVE EMISSION

part, by such a chemical source. Additional experiments utilizing more sensitive equipment are now in progress, especially searching for microwave emission from reactions occurring outside the discharge tube.

#### WATER AS A MINOR CONSTITUENT

The high atmospheric abundance of  $\text{CO}_2$  in Venus is believed to indicate an absence of water on the surface of the planet since  $\text{CO}_2$  is considered to be removed from the Earth's atmosphere, according to Urey (1952) by the equilibrium



or similar reaction with calcium silicate. In the presence of liquid water the equilibrium is shifted strongly to the right, the equilibrium  $\text{CO}_2$  pressure being about  $10^{-4}$  atmospheres. In the absence of liquid water the reaction of  $\text{CO}_2$  with silicates occurs either very slowly or not at all, thus leading to the conclusion that there is no significant amount of surface water on Venus. This assumes that the composition of the crust of Venus is about the same as that of Earth, which seems rather likely in view of the similar densities of the two planets.

In addition, the presence of large amounts of surface water would lead to significant amounts of atmospheric water vapor which would become photodissociated in the upper atmosphere and cause a build-up of atmospheric oxygen (Poole, 1941; Harteck and Jensen, 1948). Since this is not observed, any model incorporating either surface water or extensive atmospheric water vapor seems highly improbable.

It has long been known that the period of rotation of Venus is considerably longer than that of Earth and many astronomers have postulated that the period of rotation may even be equal to the period of revolution (224.7 days). The recent data obtained by Mariner II (Jet Propulsion Laboratory, 1962) on the absence of a strong magnetic field tend to support this conclusion. If it is assumed that one side

of Venus is forever turned toward the sun, then the dark side would be very cold. Thermocouple measurements of the infrared emission from the dark side of Venus give a temperature of 234°K (Sinton and Strong, 1960). At this temperature liquid CO<sub>2</sub> has a vapor pressure of about 10 atmospheres, which is just the order of magnitude assumed in many models for the Venus atmosphere. Thus, under such conditions liquid CO<sub>2</sub> may be present as lakes or even oceans.

Such a large cold area would serve as a very efficient trap for water (vapor pressure at 234°K is 0.0966 mm or about 10 parts per million of the CO<sub>2</sub>) which could be present as an ice and snow layer on the lakes or as icebergs in the CO<sub>2</sub> ocean as well as an icy coating on the solid planetary surface. Small changes in temperature will result in strong evaporation and condensation of CO<sub>2</sub>; such a situation would produce huge storms at the "twilight" zone. It must be emphasized that liquid CO<sub>2</sub> could be present only under the conditions of low surface temperatures on the dark side, high surface CO<sub>2</sub> pressure, and lack of planetary rotation. The temperature difference between the bright and dark sides may be substantially less than anticipated since on the surface of a non-rotating planet heat transfer by convection will be very efficient. In addition, it should be noted that if the planet rotates at all, even very slowly, little or no water in any form can be present, as previously discussed.

#### THE CLOUD LAYER

The composition of the clouds is a question which has been discussed for many years and still remains unsettled. Carbon suboxide has been suggested as a possible major constituent (Harteck and Groth, 1957) especially in the light of the suboxide mechanism proposed for regeneration of CO<sub>2</sub> in the presence of ionizing radiation. In the Venus atmosphere under present conditions, however, very little suboxide may be forming as discussed above. Nevertheless, it cannot be entirely excluded and further investigations should be made. However,

if Venus initially possessed a reducing atmosphere as proposed by Urey (1952) and Kuiper (1952), there may have been a time when the  $\text{CO}:\text{CO}_2$  ratio was substantially higher than at present. Under these conditions, carbon suboxide polymer should have been formed in large amounts by the action of solar radiation on the  $\text{CO}$ . Thus not only clouds, but hills and dunes of suboxide polymer may exist on Venus. It is known (Harteck and Dondes, 1963) that when carbon suboxide is formed at high temperatures ( $250\text{--}300^\circ\text{C}$ ), it appears blackened due to the presence of carbon. Thus, on the surface of Venus, over geologic ages, carbon also may have been formed by decomposition of suboxides.

Since carbon suboxides are extremely hygroscopic, small amounts of water which might be present would soon be entirely consumed. It should be noted that, in the absence of water vapor, gaseous volcanic exhalations, such as  $\text{SO}_2$  which on Earth would be washed out of the atmosphere by rain, would accumulate in the Venus atmosphere and contribute to the clouds; a fact which may explain their observed yellowish color. The possibility of the presence of gaseous sulfur has been suggested (Suess, 1963) and this may exist in thermodynamic equilibrium in the Venus atmosphere, provided that the surface temperature is reasonably high. However, if any part of the surface is cold, sulfur would condense out.

From a chemical point of view, it is interesting that the Venus atmosphere is a "neutral" atmosphere, being neither oxidizing nor reducing. This situation could be understood if the atmosphere had been initially reducing with excess  $\text{CO}$ , as mentioned above. This  $\text{CO}$  would have been transformed by ionizing radiation into carbon suboxide polymer, which would condense out, and  $\text{CO}_2$ , which would remain in the atmosphere. Other possible components of the clouds may include polymers of  $\text{CN}$  and  $\text{CNO}$  which are known to be formed by the action of ionizing radiation on  $\text{CN}$  and mixtures of  $\text{CN}$  and  $\text{O}_2$  as mentioned above. In the latter case a pale yellow polymer  $(\text{CNO})_x$  is formed. The chemical nature of these compounds is such that over geologic ages they would

tend to decompose rather than to accumulate, reaching an equilibrium steady state. It should be noted that, as previously discussed, it is extremely unlikely that the clouds can be ice crystals as Sagan (1961) has suggested.

It is also of interest to observe that Venus, like the Earth, is in a stationary state photochemically because radiation equilibrium is attained. Jupiter and other more distant planets are not in such a stationary state, however, and there will be irreversible decomposition of hydrocarbons, ammonia, etc.

#### CONCLUDING REMARKS

The atmosphere of Venus is at present a rather controversial subject. To clarify some of the aspects, a series of conditions which may exist on Venus have been simulated. Experimental measurements have been made of the absorption coefficients of  $\text{CO}_2$ ,  $\text{CO}$ ,  $\text{O}_2$ , and other gases in the region from 1850 Å to longer wavelengths. The behavior of  $\text{CO}_2$  under the impact of ionizing radiation from pressures of one atmosphere down to a few hundred microns has been studied and extrapolation made to the region of chemosphere. Reaction mechanisms which may occur and be of major importance in the Venus atmosphere have been discussed. As an alternate to the currently popular thermal origin of the observed microwave radiation emitted from Venus, a chemical source is suggested. Light-emitting chemical reactions of the type believed to occur on Venus are under study. Preliminary results show that microwave radiation in the 3 cm wavelength region is emitted from the  $\text{CO}_2$  discharge. Additional experiments in these areas are in progress.

#### ACKNOWLEDGEMENT

This work is being carried out under a grant from the National Aeronautics and Space Administration. The authors would like to thank Mr. O. L. Casstevens of Naco Electronics Corp. for supplying

microwave detection equipment. We also thank the Perkin-Elmer Corporation, Norwalk, Conn. for the use of the Model 350 absorption spectrophotometer. We would like to acknowledge the cooperation of Dr. S. Dondes who conducted the reactor irradiations. Finally, the experimental assistance of Mr. W. H. Chace and Mr. R. Waldron is acknowledged with thanks.

Rensselaer Polytechnic Institute

Troy, New York

April 19, 1963

# REFERENCES

- Adams, W. S. and T. Dunham, Absorption Bands in the Infra-red Spectrum of Venus, Publs. Astron. Soc. Pacific, 44, 243, 1932.
- Chamberlain, J. W. and G. P. Kuiper, Rotational Temperature and Phase Variation of the Carbon Dioxide Bands of Venus, Astrophys. J., 124, 399, 1956.
- Dondes, S., Radiation Chemistry of the Nitrogen-Oxygen System at Low Pressures, in The Threshold of Space, edited by M. Zelickoff, pp. 116-123, Pergamon Press, London, 1957.
- Dubrin, J., C. MacKay, and R. Wolfgang, The Reactions of Atomic Carbon with Inorganic Oxides and with Nitrogen, Presented at the 144th National Meeting, American Chemical Society, Los Angeles, California, April 1-5, 1963.
- Dunham, T., Spectroscopic Observations of the Planets at Mount Wilson, in The Atmospheres of the Earth and Planets, edited by G. P. Kuiper, pp. 288-305, Univ. of Chicago Press, 1952.
- Feast, M. W., Investigation of the Spectrum of the High-Voltage Arc in Carbon Dioxide: The Carbon Monoxide Flame Spectrum, Proc. Phys. Soc. (London), A63, 772, 1950
- Gaydon, A. G., Spectroscopy of Flames, pp. 91-111, John Wiley and Sons, Inc., New York, 1957.
- Gaydon, A. G., Spectra of Flames, Advances in Spectroscopy, II, 33, 1961.
- Groth, W., Photochemische Untersuchungen in Schumann-UV, Nr. 3, Z. Phys. Chem., B37, 307, 1937.
- Hartek, P., A Discussion of the Reactions of Nitrogen and Nitrogen Oxides in the Upper Atmosphere, in The Threshold of Space, edited by M. Zelickoff, pp. 32-39, Pergamon Press, London, 1957.
- Hartek, P. and S. Dondes, Decomposition of Carbon Dioxide by Ionizing Radiation, Part I, J. Chem. Phys., 23, 902, 1955; Part II, ibid. 26, 1727, 1957.
- Hartek, P. and S. Dondes, Radiation Chemistry of Gases, Proc. 2nd Intern. Conf. Peaceful Uses of Atomic Energy, Geneva, Vol. 29, p. 415, 1958.
- Hartek, P. and S. Dondes, Private Communication, 1963.
- Hartek, P. and W. Groth, Discussed by G. P. Kuiper, The Atmosphere and Cloud Layer of Venus, in The Threshold of Space, edited by M. Zelickoff, p. 85, Pergamon Press, London, 1957.

- Harteck, P. and J. H. D. Jensen, Über den sauerstoffgehalt der atmosphäre, Z. Naturforsch., 3a, 591, 1948.
- Harteck, P. and R. R. Reeves, Chemical Reactions in the Atmosphere, Angew. Chem. internat. Edit., 1, 25, 1962.
- Herzberg, G., Laboratory Absorption Spectra Obtained with Long Path Lengths, in The Atmospheres of the Earth and Planets, edited by G. P. Kuiper, pp. 406-416, Univ. of Chicago Press, 1952.
- Jet Propulsion Laboratory, Pasadena, California, Interplanetary Magnetic Field Measurement by Mariner II, Press Release, December 28, 1962.
- Jet Propulsion Laboratory, Pasadena, California, Press Release, February 26, 1963.
- Johnson, F. S., Editor, Satellite Environment Handbook, Stanford Univ. Press, 1961.
- Jones, D.E., The Microwave Temperature of Venus, Planetary and Space Science, 5, 155, 1961.
- Jucker, H. and B. Rideal, The Photodecomposition of Carbon Dioxide and Ammonia by Xenon 1470 Å Radiation, J. Chem. Soc., 1957, 1058, 1957.
- Kellogg, W. W. and C. Sagan, The Atmospheres of Mars and Venus, Publ. 944, Natl. Acad. Sci. - Natl. Res. Council, 1961.
- Kozyrev, N. A., Discussed by C. Sagan, The Planet Venus, Science, 133, 849, 1961.
- Kuiper, G. P., Planetary Atmospheres and Their Origin, in The Atmospheres of the Earth and Planets, edited by G. P. Kuiper, Univ. of Chicago Press, 1952.
- Kuz'min, A. D. and A. E. Salomonovich, Radio Observations of Venus in 1961, Soviet Astron. 5, 851, 1962.
- Lind, S. C., Radiation Chemistry of Gases, Reinhold Pub. Corp., 1961.
- Mahan, B. H., Photolysis of Carbon Dioxide, J. Chem. Phys., 33, 959, 1960.
- Mayer, C. H., Radio Emission of the Moon and Planets, in Planets and Satellites, edited by G. P. Kuiper and B. H. Middlehurst, pp. 455-461, Univ. of Chicago Press, 1961.

- Newkirk, G., Airglow of Venus: A Re-examination, Planetary and Space Science, 5, 163, 1961.
- Opik, E. J., The Aeolosphere and Atmosphere of Venus, J. Geophys. Res., 66, 2807, 1961
- Poole, J. H. H., The Evolution of the Atmosphere, Proc. Roy. Soc. Dublin, 22, 345, 1941.
- Prokofiev and Petrova, Reported by R. L. Newburn, Planetary Astrophysics and the Exploration of the Solar System, Paper No. 17, Proc. NASA - Univ. Conf. Sci. and Tech. of Space Exploration, NASA SP-11, Vol. 1, pp. 197-205, 1962.
- Sagan, C., The Radiation Balance of Venus, Technical Report 32-34 Jet Propulsion Laboratory, Pasadena, California, 1960.
- Sagan, C., The Planet Venus, Science, 133, 849, 1961
- Scarf, F. L., Plasma Instability and the Microwave Radiation from Venus, J. Geophys. Res., 68, 141, 1963.
- Sinton, W. M., Infrared Spectroscopy of Planets and Stars, Applied Optics, 1, 105, 1962.
- Sinton, W. M. and J. Strong, Radiometric Observations of Venus, Astrophys. J., 131, 470, 1960
- Spinrad, H., Spectroscopic Temperature and Pressure Measurements in The Venus Atmosphere, Publ. Astron. Soc. Pacific, 74, 187, 1962.
- Strong, J., Reported by C. Sagan, The Planet Venus, Science, 133, 849, 1961.
- Suess, H., Private Communication, 1963.
- Tolbert, C. W. and A. W. Straiton, A Consideration of Microwave Radiation Associated with Particles in the Atmosphere of Venus, J. Geophys. Res., 67, 1741, 1962.
- Urey, H. C., The Planets: Their Origin and Development, Yale Univ. Press, New Haven, 1952.
- von Weyssenhoff, H., S. Dondes, and P. Harteck, The Reaction of Carbon Suboxide with Oxygen Atoms, J. Am. Chem. Soc., 84, 1526, 1962.

von Weyssenhoff, H., P. Hardeck, and S. Dondes, To be published,  
1963.

Watanabe, K., M. Zelikoff, and E. C. Y. Inn, Absorption Coefficients  
of Several Atmospheric Gases, AFCRC Tech. Rpt. 53-23, 1953.

